

201

Minnesota Women's Health Report Card

A Snapshot of Minnesotan Women's Health (2016-2017)

> A publication of the Center for Leadership Education in Maternal and Child Public Health, University of Minnesota School of Public Health



Find supporting data, citations and other information at z.umn.edu/mnwhrc18.



DEMOGRAPHICS

2016-2017

Minnesota's (MN) population is gradually increasing. From 2016 to 2050, MN's population is projected to increase by 15%, from 5.53 million to 6.36 million. Currently, female population growth is slightly slower than male population growth. It's expected that the female growth rate is projected to remain lower than male's population past 2020.

In 2017, the total female population represented 50% (2,804,325) of the state's population. The number of females ages 15-44 was 1,066,806, about 38% of all MN women, representing a 1% change from 2016 to 2017. Women over 80 accounted for 60% of the population in rural MN, 62% in small town MN, 64% in large town MN and 63% in urban MN.

FEMALES' LIFE **EXPECTANCY YEARS**

Ethnicity

| 84.5% | White |
|-------|--|
| 6.4% | Black or African American |
| 1.3% | American Indian |
| 5.2% | Asian |
| 0.1% | Native Hawaiian and Other Pacific Islander |
| 2.5% | Two or More Races |

Age

| 6% | 15-19 174,958 | | | | | | |
|-----|---------------|---------|---------|--|--|--|--|
| 13% | 20-29 | 360,094 | L | | | | |
| 13% | 30-39 | 374,031 | | | | | |
| 25% | 40-59 | | 712,133 | | | | |
| 23% | 60+ | | 648,947 | | | | |
| | | | | | | | |

Population Count

Key

Urban: 50.000+ residents

Large towns: 10,000-49,999 residents Small towns: 2,500-9,999 residents

Rural: Primary commuting flow is outside of urban areas/clusters

Source: Greater Minnesota: Refined & Revisited report

This Report Card is produced using a variety of data sources (on page 8) ranging from 2011-2018. Much of the data used for this report card were reported in 2017; if different, the data collection year is indicated.

BARRIERS TO HEALTH

Minority or

non-white women

are three times

more likely to be uninsured than white women.

7%
FEMALES
AGES 18-64 IN MN
ARE UNINSURED

UNINSURED RATE BY RACE:

WHITE

4%

NON-WHITE

12%

2017

In 2017 there were an estimated 2,199,595 females ages 16+ in MN, 66% of whom were in the labor force. Females account for 42% of total uninsured population in MN. Minority or non-white women are three times more likely to be uninsured than white women.

Between 2015 and 2018, the amount of homelessness among adults ages 25–54 rose. In 2017, 2,875 females ages 18–54 were experiencing homelessness.

OF HOMELESS WOMEN ARE 25-54 YEARS OLD

| HOMELESSNESS AMONG MN WOMEN |
|-----------------------------|
| PER AGE GROUP IN 2018 |

| AGE GROUP | ADULTS FEMALE |
|-----------|---------------|
| 18-21 | 429 |
| 22-24 | 299 |
| 25-54 | 2,147 |
| 55+ | 264 |

77%
OF WOMEN ARE BELOW THE POVERTY LINE

LABOR FORCE PARTICIPATION BY RACE (DRODODTION)

| RACE | PROPORTION |
|------------------------------------|-------------------|
| White Non-Hispanic | 81% |
| Asian | 70% |
| Black | 71% |
| American Indian | 65% |
| Latino | 73% |
| All Women | 79% |
| Black American Indian Latino | 71% 65% 73% |



PREVENTIVE HEALTH

2016

Overall, MN women are exceeding the Healthy People 2020 (HP2020) targets for several preventive health targets. Women ages 50-74 have a slightly higher mammogram rate (82%) compared to the HP2020 target (81%). Black women have the highest cancer screening rate (91%), followed by screening for white women (83%). However, the sample size used for black women and non-white Hispanics in the Behavioral Risk Factor Surveillance System (BRFSS) is too small to reveal the right coverage of breast cancer screening among this groups giving the relatively high risk of this type of cancer among black women.



| 72% | Women ages 50-75 who have ever had a colonoscopy (in the past 10 years) |
|-----|--|
| 82% | Women ages 50-74 who have had a mammogram in last 2 years |
| 78% | Women who have visited the dentist/dental clinic in the past year for any reason |
| 82% | Women 21-65 who have had a pap test within the past 3 years |
| 77% | Women who had a routine checkup in the past year |
| 24% | Women were obese pre-pregnancy |
| 76% | Women who participated in physical activities in the past month (2017) |
| 47% | Women who exercise 3+ days a week |
| 72% | Women with a preventive medical visit (in the past year) |
| 59% | Rate of HPV vaccination among young adults (ages 13-17) >=1 HPV vaccine |
| 46% | Rate of women who are up-to-date on HPV vaccine |
| | ■ Above HP2020 target ■ Below HP2020 target |

The health indicators featured in this section mirror CDC BRFSS indicators. Not all indicators are compared to the HP2020 targets because the HP2020 targets and BRFSS indicators do not always exactly match. Visit cdc.gov/brfss and healthypeople.gov for more information.




PERINATAL HEALTH AND REPRODUCTIVE HEALTH

2017

In 2017, there were a total of 78,227 pregnancies in MN. The pregnancy rate was 73.3 per 1,000 women ages 15-44. The state counted a total of 68,603 live births (all singleton and multigestation included), of which 9% were premature (less than 37 weeks of gestation) and 7% were low birth weight (less than 2,500 grams/5 pounds, 8 ounces).



Singleton and Multigestational Births





Top Four Countries of Origin for Women Giving Birth in MN





OF MN WOMEN SMOKE **BEFORE OR DURING PREGNANCY**



Live Births by Race and Ethnicity of Mother n=69,746

| NON HISPANIC | | | | | HISPANIC | TOTAL |
|--------------|-------------------|--------------------|---------------------------|-------------------|----------|--------|
| White | Black American | American Indian | Asian/Pacific Islander | Other/ unknown | | |
| 47,890 | 8,664 | 1,197 | 5,492 | 466 | 4,894 | 68,603 |

OF YOUNG WOMEN OF REPRODUCTIVE AGE ADOPT CONTINUOUS-USE LONG-ACTING REVERSIBLE **CONTRACEPTIVES (LARCS)** IN MN HEALTH CARE PROGRAMS (MEDICAID)



PERINATAL HEALTH AND REPRODUCTIVE HEALTH (Continued)



Postpartum Depression

11%

OF MOTHERS WERE TOLD BY A PROVIDER THEY HAD DEPRESSION BEFORE PREGNANCY

13%

OF MOTHERS SELF-REPORTED POSTPARTUM DEPRESSION SYMPTOMS



Gestational Diabetes and Gestational Hypertension

IN 2015,


6%

OF MN BIRTHS INVOLVED MOTHERS
WHO HAD GESTATIONAL DIABETES;
this does not include women who had type 1
or type 2 diabetes before pregnancy.



STIs reported cases

These numbers are reported by MDH and include the sexually transmitted infections (STIs) reported in different services throughout the state. It should be noted that the All Payers Claim Database would be a better resource to capture the number of individuals tested.



Gonorrhea

In the 20–24 age group, males have a slightly higher rate of gonorrhea cases per 100,000 people compared to female cases of gonorrhea, with 500 and 478 cases per 100,000 people respectively.

Females age 20–24 have the highest rate of chlamydia in the state.

(~3,400 cases per 100,000 person)

Chlamydia

In general, the rate of chlamydia in MN reached an all time high at 444 per 100,000 people. This is an increase of 4% from 2016. The rate of gonorrhea in MN increased 28% to 123 per 100,000 compared to 96 per 100,000 in 2016. Females age 20–24 have the highest rate of chlamydia in the state (~3,400 cases per 100,000 person).

Syphilis

The lowest case of early syphilis among women occurred in 2007, with two reported cases. This number has been steadily increasing since then and reached 91 cases in 2017. The rate of primary and secondary syphilis is at 5.5 per 100,000. This is a decrease of 5% from 2016.

THE FOLLOWING SHARE THE LARGEST PROPORTIONS OF THE NUMBER OF WOMEN INFECTED WITH SYPHILIS

AMERICAN INDIAN

35%

WHITE NON-HISPANIC

29%

BLACK NON-HISPANIC

23%

Breast cancer

accounts for 37%

of the top ten cancers that occurred among MN women in 2015.



2015

Cancer is the leading cause of death in MN among men and women. When all cancers are combined into one group, cancer is the 7th most frequent chronic disease. Breast cancer accounts for 37% of the top ten cancers that occurred among MN women in 2015. In general, lung and bronchus cancer constitute the leading cause of cancer deaths, with the second highest rate of incidence but the number one death rate.

Rate of Top Ten Cancers among Women

| CANCER TYPE | AGE ADJUSTED RATE |
|--------------------------|-------------------|
| Female Breast | 134.5 |
| Lung and Bronchus | 53.3 |
| Colon and Rectum* | 34.2 |
| Corpus and Uterus NOS | 31.7 |
| Melanomas of the Skin | 28.2 |
| Thyroid* | 18.7 |
| Non-Hodgkin Lymphoma | 17.4 |
| Leukemias | 12.3 |
| Pancreas* | 12.2 |
| Kidney* and Renal Pelvis | 12 |

Per 100,000 women

Mortality Rate/Type of Cancer

| CANCER TYPE | AGE ADJUSTED RATE |
|------------------------------------|-------------------|
| Lung and Bronchus | 33.3 |
| Female Breast* | 19.2 |
| Colon and Rectum | 11.1 |
| Pancreas | 10 |
| Ovary* | 6.4 |
| Leukemias | 4.6 |
| Non-Hodgkin Lymphoma | 4.4 |
| Corpus and Uterus NOS*+ | 4.4 |
| Brain and Other Nervous System | 4.1 |
| Liver* and Intrahepatic Bile Duct* | 3.1 |

Per 100,000 women



COMMUNITY AND ENVIRONMENTAL HEALTH

2015-2018



INTIMATE PARTNER VIOLENCE (IPV)

In 2015, 2% of pregnant women experienced IPV during the 12 months before pregnancy and 2% experienced IPV during pregnancy.

HOMICIDES

In 2017, there were 33 reported cases of female homicide victims. 1.5x

SUICIDE RATES

The rate is 6 per 100,000. Females are 1.5 times more likely to die by suicide than homicide in MN.

7%

INCARCERATION RATES

There are 649 female adults in prison (7% of the total prison population) in MN (As of January 1, 2019).

⁺Not otherwise specified

^{*}Cancer types associated with obesity



2012-2017

MN women have a slightly higher rate of chronic diseases (38%) compared to men (33%). In addition, women had a higher rate of certain chronic conditions such as high blood pressure, asthma and rheumatoid arthritis.



41%

Women are at a

recommended weight

31% of women are overweight

26% of women are obese

5%
of women
are underweight
or unknown

% of Women Ever Told By A Health Professional They have had a:



HEART ATTACK
(also called
myocardial infarction)



ANGINA OR CORONARY HEART DISEASE



STROKE
(also called a
cerebrovascular accident)

- 7% Women ever told by a health professional they have had diabetes
- 24% Women ever told they had high blood pressure
- 27% Women ever told by a health professional they have high cholesterol
- 9% Women currently with asthma
- 5% Women ever told they had some form of COPD, emphysema, or chronic bronchitis
- 23% Women ever told they had some form of arthritis
- 3% Women ever told by a health professional the have had kidney disease
- 41% Women who are at a recommended weight
- 31% Women who are overweight
- 26% Women who are obese
- 5% Women who are underweight or unknown

RATE OF WOMEN WITH THE FOLLOWING CHRONIC CONDITIONS:

- 214.3 High Blood Pressure
- 147.0 High Cholesterol
- 124.1 Asthma
- 62.1 Diabetes
- 74.9 Depression
- 25.3 Congestive Heart Failure
- 14.9 Ischemic Heart Disease
- 12.9 Chronic Kidney Disease
- 12.6 Chronic Obstructive Pulmonary Disease (COPD)
- 15.1 Rheumatoid Arthritis
- 378.3 Any Condition

Rate per 1,000 Minnesotans



MENTAL HEALTH AND SUBSTANCE ABUSE

2011-2016

Women constitute 55% of the total clients served by the state's mental health agencies, in community settings and at state hospitals.

The penetration rate (the rate of individuals who utilize mental health services) for females is 53.7 per 1,000. This rate is higher than males which is 44 per 1,000.



TOTAL CLIENTS SERVED BY THE STATE'S MENTAL HEALTH AGENCIES, IN COMMUNITY **SETTINGS AND AT** STATE HOSPITALS

An Adverse Childhood Experience (ACE) is described as a traumatic experience in a person's life which occurs before the age of 18 that the person recalls as an adult.

Adverse Childhood Experience (ACE)

We are reporting ACEs in this report because they are associated with poor physical and mental health, chronic disease, lower educational achievement, lower economic success and impaired social success in adulthood. ACEs are strongly associated with indicators of mental health issues later in life, and the higher the ACE score, the more likely adults are to report depression or anxiety in adulthood. The MN BRFSS conducted in 2011 present the most recent data for adults. It shows that, compared to men, women more often report experiencing sexual abuse and witnessing domestic violence, mental illness and problems related to drinking in the household.



57% WOMEN REPORTED EXPERIENCING ONE OR MORE ACES IN CHILDHOOD IN 2011

Women 18 to 65 are currently smokers



24% WOMEN REPORTED **EXPERIENCING** THREE OR MORE ACES IN CHILDHOOD



For women, having 4+ drinks in one occasion is classified as binge drinking.

Excessive alcohol use

Excessive alcohol use can result in a variety of harms such as poor birth outcomes, cancer, heart disease, motor vehicle injuries, and more. For women, having 4+ drinks in one occasion is classified as binge drinking. Any drinking by pregnant women or people under the age of 21, or women having 8+ drinks in a week, is classified as heavy drinking.

| 1370 | Tromen to 65 are currently smokers |
|------|---|
| 60% | Women 18 and older reported alcohol use in 2016 |
| 24% | Women ages 18-44 were more likely to say they binge drink than those older than 44 years (10%) |
| 18% | Pregnant women said they drank alcohol in the last month |
| >5% | Pregnant women said they binge drank in the past month. |
| 3% | Women reporting any use of prescription drugs not prescribed for them by their doctor within the past 12 Months |

2018 Minnesota Women's Health Report Card: A Snapshot of Minnesotan Women's Health (2016-2017)

A publication of the Center for Leadership Education in Maternal and Child Public Health, University of Minnesota School of Public Health

About:

Working closely with colleagues and the Minnesota Department of Health's (MDH) Center for Health Statistics, Child and Family Health Division, and Health Promotion and Chronic Disease Division, we have carefully designed the Minnesota Women's Health Report Card (MN-WHRC) to help visualize and reflect the health indicators specific to our state's unique demographics. The MN-WHRC will be produced every other year as new data becomes available, and with the intent of providing a comprehensive review and comparison across the years. These data reflect the daily lived experiences of women's lives, and can be used to inform Minnesota's practice and policies, thus offering the opportunity of improving the near- and long-term health outcomes for women.

About the Data Sources:

The Report Card is produced using a variety of data sources ranging from 2011-2018. Much of the data used for this report card were reported in 2017; however, if different, the data collection year is indicated next to each heading or topical areas. Data collection methods may also vary by year, and sample sizes can differ for each data reporting system. Percentages have been rounded for simplicity, with full percentages available in the online version of the MN-WHRC.

Unless otherwise indicated, all data are for women ages 18+. One of the limitations of the data sources is that some data reported were not broken down by gender and race/ethnicity. We share data by race and ethnicity for better targeting of resources and interventions toward populations in need. The terms "female" and "woman/women" are used interchangeably. While they are not inclusive of all gender identities, they are used because they reflect data source language.

Acknowledgements

Acknowledgements, full data citations and additional details can be found at **z.umn.edu/mnwhrc18**. This report card was modeled after North Carolina's (NC) Center for Women's Health Research (CWHR), which has been reporting on the health status of NC's women for several years. Their report cards and corresponding website inspired us to create the same for our state. Thank you.

About the Center for Leadership Education in Maternal and Child Public Health

For six decades, the Center for Leadership Education in Maternal and Child Public Health has worked to improve the health status of MN's women, children, adolescents and their families. The Center's work is rooted in principles of social justice and focuses on skills development through continuing professional education, consultation and technical assistance to community-based organizations and public health and social service agencies. Students pursuing our Master's degree in Public Health receive training that focuses on public health skills development and maternal and child health (MCH) content.

CONTACT:

epi.umn.edu/mch mch@umn.edu









The Center for Leadership Education in Maternal and Child Public Health is supported by the Health Resources and Services Administration (HRSA) of the U.S. Department of Health and Human Services (HHS) under grant number T76MC00005-59-00 for Leadership Education in Maternal and Child Public Health in the amount of \$1,750,000. This information or content and conclusions of our outreach products are those of the authors and should not be construed as the official position or policy of, nor should any endorsements be inferred by HRSA, HHS or the U.S. Government.

Design and Printing: University of Minnesota Printing Services

